### **Postoperative Sepsis Rate**

# Patient Safety Indicators #13 Technical Specifications

Provider-Level Indicator
AHRQ Quality Indicators, Version 4.3, August 2011

#### **Numerator**

Discharges among cases meeting the inclusion and exclusion rules for the denominator with ICD-9-CM code for sepsis in any secondary diagnosis field.

#### ICD-9-CM Sepsis diagnosis codes:

in ocposo diagnosio codes.			
STREPTOCOCCAL SEPTICEMIA	Septice	Septicemia due to:	
STAPHYLOCOCCAL SEPTICEMIA	03840	GRAM-NEGATIVE ORGANISM,	
STAPHYLOCOCCAL SEPTICEMIA,		UNSPECIFIED	
UNSPECIFIED	03841	HEMOPHILUS INFLUENZAE	
METHICILLIN SUSCEPTIBLE	03842	ESCHERICHIA COLI	
STAPHYLOCOCCUS AUREUS	03843	PSEUDOMONAS	
SEPTICEMIA (OCT08)	03844	SERRATIA	
METHICILLIN RESISTANT	03849	SEPTICEMIA DUE TO OTHER GRAM-	
STAPHYLOCOCCUS AUREUS		NEGATIVE ORGANISMS	
SEPTICEMIA (OCT08)	0388	OTHER SPECIFIED SEPTICEMIAS	
OTHER STAPHYLOCOCCAL	0389	UNSPECIFIED SEPTICEMIA	
SEPTICEMIA	99591	SYSTEMIC INFLAMMATORY RESPONSE	
PNEUMOCOCCAL SEPTICEMIA		SYNDROME DUE TO INFECTIOUS	
(STREPTOCOCCUS PNEUMONIAE		PROCESS W/O ORGAN DYSFUNCTION	
SEPTICEMIA)	99592	SYSTEMIC INFLAMMATORY RESPONSE	
SEPTICEMIA DUE TO ANAEROBES		SYNDROME DUE TO INFECTIOUS	
SEPTIC SHOCK		PROCESS W/ ORGAN DYSFUNCTION	
SHOCK WITHOUT MENTION OF			
TRAUMA, OTHER			
POSTOPERATIVE SHOCK			
	STREPTOCOCCAL SEPTICEMIA STAPHYLOCOCCAL SEPTICEMIA STAPHYLOCOCCAL SEPTICEMIA, UNSPECIFIED METHICILLIN SUSCEPTIBLE STAPHYLOCOCCUS AUREUS SEPTICEMIA (OCT08) METHICILLIN RESISTANT STAPHYLOCOCCUS AUREUS SEPTICEMIA (OCT08) OTHER STAPHYLOCOCCAL SEPTICEMIA PNEUMOCOCCAL SEPTICEMIA (STREPTOCOCCUS PNEUMONIAE SEPTICEMIA) SEPTICEMIA DUE TO ANAEROBES SEPTIC SHOCK SHOCK WITHOUT MENTION OF TRAUMA, OTHER	STREPTOCOCCAL SEPTICEMIA STAPHYLOCOCCAL SEPTICEMIA STAPHYLOCOCCAL SEPTICEMIA, UNSPECIFIED METHICILLIN SUSCEPTIBLE STAPHYLOCOCCUS AUREUS STAPHYLOCOCCUS AUREUS SEPTICEMIA (OCT08) METHICILLIN RESISTANT STAPHYLOCOCCUS AUREUS SEPTICEMIA (OCT08) OTHER STAPHYLOCOCCAL SEPTICEMIA SEPTICEMIA PNEUMOCOCCAL SEPTICEMIA (STREPTOCOCCUS PNEUMONIAE SEPTICEMIA) SEPTICEMIA SEPTICE	

<sup>\*</sup>Not valid for discharges effective October 1, 2004

#### **Denominator**

All elective\* surgical discharges age 18 and older defined by specific DRGs or MS-DRGs and an ICD-9-CM code for an operating room procedure.

\*Elective - SID Admission type # is recorded as elective (Admission Type = 3) See *Patient Safety Indicators Appendices*:

- Appendix A Operating Room Procedure Codes
- Appendix D Surgical Discharge DRGs
- Appendix E Surgical Discharge MS-DRGs

#### Exclude cases:

- with principal diagnosis of sepsis or secondary diagnosis present on admission
- with principal diagnosis of infection or secondary diagnosis present on admission
- with any code for immunocompromised state or cancer

## AHRQ QI, Patient Safety Indicators #13, Technical Specifications, Postoperative Sepsis Rate www.qualityindicators.ahrq.gov

- MDC 14 (pregnancy, childbirth, and puerperium)
- with length of stay of less than 4 days
- with missing gender (SEX=missing), age (AGE=missing), quarter (DQTR=missing), year (YEAR=missing) or principal diagnosis (DX1=missing)

See Patient Safety Indicators Appendices:

- Appendix F Infection Diagnosis Codes
- Appendix H Cancer Diagnosis Codes
- Appendix I Immunocompromised State Diagnosis and Procedure Codes